

## Article

# Deletion of the epigenetic regulator GcnE in *Aspergillus niger* FGSC A1279 activates the production of multiple polyketide metabolites

Wang, Bin, Li, Xuejie, Yu, Dou, Chen, Xiaoyi, Tabudravu, Jioji, Deng, Hai and Pan, Li

Available at <http://clock.uclan.ac.uk/24879/>

*Wang, Bin, Li, Xuejie, Yu, Dou, Chen, Xiaoyi, Tabudravu, Jioji ORCID: 0000-0002-6930-6572, Deng, Hai and Pan, Li (2018) Deletion of the epigenetic regulator GcnE in Aspergillus niger FGSC A1279 activates the production of multiple polyketide metabolites. Microbiological Research, 217 . pp. 101-107. ISSN 0944-5013*

It is advisable to refer to the publisher's version if you intend to cite from the work.

<http://dx.doi.org/10.1016/j.micres.2018.10.004>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the [policies](#) page.

**Deletion of the epigenetic regulator GcnE in *Aspergillus niger* FGSC  
A1279 activates the production of multiple polyketide metabolites**

Bin Wang<sup>a,b,\*</sup>, Xuejie Li<sup>a,b,\*</sup>, Dou Yu<sup>a,b</sup>, Xiaoyi Chen<sup>a,b</sup>, Jioji Tabudravu<sup>c,d</sup>, Hai Deng<sup>a,b,c,#</sup>, Li  
Pan<sup>a,b,#</sup>

<sup>a</sup>School of Biology and Biological Engineering, South China University of Technology,  
Guangzhou Higher Education Mega Center, Guangzhou, 510006, China

<sup>b</sup>Guangdong Provincial Key Laboratory of Fermentation and Enzyme Engineering, South  
China University of Technology, Guangzhou Higher Education Mega Center, Guangzhou  
510006, China

<sup>c</sup>Marine Biodiscovery Centre, Department of Chemistry, University of Aberdeen, Meston  
Walk, Aberdeen AB24 3UE, Scotland, UK

<sup>d</sup>School of Forensics and Applied Sciences, Faculty of Science & Technology, University of  
Central Lancashire, Preston, Lancashire, PR1 2HE, UK

**\* These authors contributed equally to this work.**

**# Corresponding authors**

Hai Deng: h.deng@abdn.ac.uk

Li Pan: btlipan@scut.edu.cn

**Declarations of interest:** none.

## ABSTRACT

Epigenetic modification is an important regulatory mechanism in the biosynthesis of secondary metabolites in *Aspergillus* species, which have been considered to be the treasure trove of new bioactive secondary metabolites. In this study, we reported that deletion of the epigenetic regulator *gcnE*, a histone acetyltransferase in the SAGA/ADA complex, resulted in the production of 12 polyketide secondary metabolites in *A. niger* FGSC A1279, which was previously not known to produce toxins or secondary metabolites. Chemical workup and structural elucidation by 1D/2D NMR and high resolution electrospray ionization mass (HR-ESIMS) yielded the novel compound nigerpyrone (**1**) and five known compounds: carbonarone A (**2**), pestalamide A (**3**), funalenone (**4**), aurasperone E (**5**), and aurasperone A (**6**). Based on chemical information and the literature, the biosynthetic gene clusters of funalenone (**4**), aurasperone E (**5**), and aurasperone A (**6**) were located on chromosomes of *A. niger* FGSC A1279. This study found that inactivation of GcnE activated the production of secondary metabolites in *A. niger*. The biosynthetic pathway for nigerpyrone and its derivatives was identified and characterized via gene knockout and complementation experiments. A biosynthetic model of this group of pyran-based fungal metabolites was proposed.

## Keywords

Histone acetyltransferase GcnE, Secondary metabolite, Epigenetic regulator, Polyketide, Nigerpyrone

## 1. Introduction

The genus *Aspergillus* is a well-known producer of secondary metabolites (SMs) including polyketides, non-ribosomal peptides, indole terpenes and terpenes (Varga et al., 2003; Bok et al., 2006; Sanchez et al., 2012), which are important resources for new drug discovery (Rossano et al., 1999; Fisch et al., 2009). For example, bioinformatic analysis of four

available genomes of *A. niger* species (NRRL3, ATCC 9029) (Baker, 2006), ATCC1015 (Andersen et al., 2011), CBS513.88 (Pel et al., 2007), and SH2 (Yin et al., 2014), indicates that *A. niger* encodes at least 81 putative SM biosynthetic gene clusters (Inglis et al., 2013) for 42 polyketides (PKS), 33 non-ribosomal peptides (NRPS), four PKS-NRPS hybrids and two indole alkaloids. The large number of putative SM biosynthetic gene clusters suggests that *A. niger* has the potential to produce abundant SM products. However, most of these putative gene clusters are silent in standard laboratory cultivation conditions (Fisch et al., 2009), and only limited SM products have been characterized from *A. niger* (Tanaka et al., 1966; Hiort et al., 2004; Serra et al., 2005; Nielsen et al., 2009; Sorensen et al., 2009; Chiang et al., 2011).

Epigenetic modification is an important regulatory mechanism in the biosynthesis of SM products in *Aspergilli* (Fisch et al., 2009), which could activate SM gene clusters and induce new SM products by changing the chromosomal region of SM gene clusters from a heterochromatic to a euchromatic state via histone demethylation and acetylation (Bayram et al., 2008; Fisch et al., 2009). For example, 10 novel SM compounds are induced by adding the epigenetic modification reagent 5-azacytidine (5-AZA) to *A. niger* ATCC1015 cultures (Fisch et al., 2009).

Spt-Ada-Gcn5-acetyltransferase (SAGA/ADA) complex is an epigenetic regulator in fungal secondary metabolism (Wu and Yu, 2015). GcnE (the Gcn5 homolog in *Aspergilli*) is a histone acetyltransferase (HAT) in SAGA/ADA complex that might regulate secondary metabolism by histone modification (Baker and Grant, 2007; Brakhage, 2013). Previous work illustrated that GcnE participates in increasing the acetylation level of histone H3 lysine K9 in *Aspergilli* (Reyes-Dominguez et al., 2008), and affects the biosynthesis of secondary metabolites. For example, *A. flavus*  $\Delta gcnE$  mutant does not produce aflatoxin (Lan et al.,

2016). Deletion of the *gcnE* gene in *A. nidulans* decreases the production of orsellinic acid, sterigmatocystin, penicillin and terrequinone (Nutzmann et al., 2011).

*A. niger* strain FGSC A1279 (*kusA::DR-amdS-DR, pyrG<sup>-</sup>*), a derivative of industrial strain NRRL3 (ATCC9029) that is used for gluconic acid production (Baker, 2006; Yuan et al., 2008; Carvalho et al., 2010; Arentshorst et al., 2015), is an SM-silent strain and has zero level of SM production. The low background of SM production in FGSC A1279 makes it ideal for investigating the role of epigenetic regulators on SM production. Moreover, genome sequences of NRRL3 (ATCC 9029) (Baker, 2006) and FGSC A1279 (Wang et al., 2017) are available for detailed gene information of the host strain.

In this study, we reported that deletion of the epigenetic regulator *gcnE*, a histone acetyltransferase in the SAGA/ADA complex, resulted in the production of 12 polyketide secondary metabolites in *A. niger* FGSC A1279. Chemical workup and structural elucidation by 1D/2D NMR and high resolution electrospray ionization mass (HR-ESIMS) yielded the novel compound nigerpyrone (**1**) and five known compounds: carbonarone A (**2**), pestalamide A (**3**), funalenone (**4**), aurasperone E (**5**), and aurasperone A (**6**). Based on chemical information and literature searches, the biosynthetic gene clusters of funalenone (**4**), aurasperone E (**5**), and aurasperone A (**6**) were located on the chromosomes of *A. niger* FGSC A1279. This study showed that inactivation of GcnE activated the production of toxins and metabolites in *A. niger*. This is distinct from the role of GcnE in previous studies, suggesting that the role of GcnE may have diverse effects on regulating SMs among different *Aspergilli*. The biosynthetic pathway of nigerpyrone and its derivatives was identified via gene knockout and complementation experiments, and a biosynthetic model of this group of pyran-based fungal metabolites was proposed.

## **2. Material and methods**

### **2.1. Strains and culture conditions**

All strains used in this study are in Table S1. Strains were maintained on potato dextrose agar (PDA) medium (20 g dextrose, 15 g agar, infusion from 200 g potatoes per 1 L medium, pH6.0) at 30 °C for spore harvest. For genetic transformation, CD medium (0.3% NaNO<sub>3</sub>, 0.2% KCl, 0.05% MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.001% FeSO<sub>4</sub>·7H<sub>2</sub>O, 0.1% K<sub>2</sub>HPO<sub>4</sub>, 2.0% glucose, pH5.5) was used. WATM, YES, and CYA media were used for SM profile analysis. WATM medium was composed of 2.0 g/l yeast extract, 3.0 g/l peptone, 2.0 g/l dextrose, 30.0 g/l sucrose, 5.0 g/l corn steep solids, 2.0 g/l NaNO<sub>3</sub>, 1.0 g/l K<sub>2</sub>HPO<sub>4</sub>·3H<sub>2</sub>O, 0.5 g/l MgSO<sub>4</sub>, 0.2 g/l KCl, 0.01 g/l FeSO<sub>4</sub>·7H<sub>2</sub>O, pH7.0. YES medium was composed of 20 g/l yeast extract, 150 g/l sucrose, pH 6.0. CYA medium was prepared on the basis of CD medium by adding 5 g/l yeast extract and 1 ml/l trace metal solution (1% ZnSO<sub>4</sub>·7H<sub>2</sub>O, 0.5% CuSO<sub>4</sub>·5H<sub>2</sub>O). For solid cultivation, 15 g/l agar was added.

An *A. niger gcnE* deletion mutant ( $\Delta gcnE$ ) was constructed according to the method developed by Szewczyk *et al.* (Szewczyk *et al.*, 2006), based on homologous recombination using a PCR amplified deletion cassette containing the upstream flanking arm (1.4 kb) of *gcnE*, the selection marker *pyrG* (orotidine-5'-phosphate decarboxylase) and the downstream flanking arm of *gcnE*. The *pyrG* gene was used for auxotrophic selection. Uridine prototrophic transformants were selected and confirmed by PCR amplification (Fig. S1). Primers for deletion cassette construction and transformant identification are listed in Table S2. When required, 10 mM uridine was added to medium to maintain auxotrophy. The SM backbone gene *epaA* was knocked out using the same method (Table S3).

## **2.2. Cultivation of *A. niger* $\Delta gcnE$ mutant for secondary metabolite collection**

To collect secondary metabolites, the *A. niger*  $\Delta gcnE$  mutant was cultivated for 7 days at 25 °C in the dark on 10 Petri dishes with WATM medium. Cultures were extracted with ethyl acetate (EtOAc) plus 1% formic acid for 24 hours. The crude extract was filtered and dried on a rotary evaporator at 38 °C.

### **2.3. SPE separation of *A. niger* $\Delta$ gcnE crude SM extract**

A C18 solid-phase extraction (SPE) column (SPE C18-E Giga Tube 20 g/60 ml, Strata, Phenomenex, USA) was used to separate *A. niger*  $\Delta$ gcnE crude extracts. The SPE stationary phase was conditioned by sequential washing with 60 ml 100% methanol and 240 ml 100% deionized water. *A. niger*  $\Delta$ gcnE crude extract (4.0 g) was chromatographed with four subfractions (480 ml each) generated stepwise from deionized water to methanol: 25% methanol, 50% methanol, 100% methanol, and 100% methanol plus 0.1% trifluoroacetic acid (TFA). Subfraction 2 (50% methanol, 50 mg) and 3 (100% methanol, 410 mg) were further purified by semi-preparative HPLC (semi-prep HPLC).

### **2.4. Semi-prep HPLC fractionation**

Semi-prep HPLC was used to purify subfractions obtained by SPE separation. A semi-prep HPLC column ACE C18-HL (250 mm  $\times$  10 mm i.d., Advanced Chromatography Technologies, UK) was connected to an Agilent 1200 series binary pump and monitored by an Agilent photodiode array detector. Detection was at 230, 254, 280, and 410 nm. Flow rate was 1.5-2.0 ml/min. Solvent A consists of 95% H<sub>2</sub>O, 5% methanol, and 0.05% TFA. Solvent B was 100% acetonitrile. Separations were done at room temperature (20-25°C). Optimal separation conditions were determined using analytical systems and then conditions were adapted to the semi-preparative scale.

### **2.5. LC/HR-ESIMS assay of *A. niger* secondary metabolites**

For LC/HR-ESIMS analysis, each semi-prep HPLC subfraction was dissolved in MeOH:Milli-Q H<sub>2</sub>O (9:1), and centrifuged at 10,000 rpm for 10 min. LC/HR-ESIMS analysis was conducted as follows. High resolution mass spectrometric data were obtained using an LTQ XL/LTQ Orbitrap Discovery MS system (Thermo Fisher Scientific, Waltham, MA, USA) coupled to a Thermo Instruments HPLC system (Accela PDA detector, Accela PDA autosampler, and Accela pump, C18 SunFire 150  $\times$  4.6 mm Waters). Conditions used were:

capillary voltage 45 V, capillary temperature 320°C, auxiliary gas flow rate 10-20 arbitrary units, sheath gas flow rate 40-50 arbitrary units, spray voltage 4.5 kV, and mass range 100-2000 amu (maximum resolution 30,000). For LC/HR-ESIMS, a Waters SunFire C18 analytical HPLC column (5  $\mu$ m, 4.6  $\times$  150 mm) was used with mobile phase of 0-100% MeOH over 30 min at a flow rate of 1 ml/min. LC/HR-ESIMS data were analyzed by Xcalibur software followed by dereplication using Antibase2012 database.

## **2.6. NMR characterization of *A. niger* secondary metabolites**

NMR spectra were used to elucidate the structure of *A. niger* secondary metabolites. <sup>1</sup>H NMR, <sup>13</sup>C NMR, COSY, HSQC and HMBC spectra were recorded on a Bruker Daltonics Advance III HD 400 NMR spectrometer, using deuterated solvent methanol-d<sub>4</sub> (Cambridge Isotopes Laboratories, USA). Chemical shift values were reported in parts per million (ppm) with tetramethylsilane (TMS) as the external standard. Coupling constants were recorded in Hertz (Hz). When higher field was needed, NMR spectra were recorded on a Bruker Daltonics Advance III HD 600 NMR. NMR data were analyzed using MestReNova 9.0.1 software and ACD/I-Lab (<http://ilab.cds.rsc.org/?cdsrdr=1>).

## **2.7. Characterization of putative biosynthetic gene clusters for *A. niger* secondary metabolites by gene knockout**

Gene knockout of putative cluster genes for identified *A. niger* secondary metabolites was done based on homologous recombination using a PCR-amplified deletion cassette containing the upstream and downstream flanking arms of the target gene and the *ptrA* (pyrithiamine (PT) resistance) selection marker. Primers for constructing deletion cassettes are listed in Table S3. 0.5  $\mu$ g/ml pyrithiamine (PT) was used to screen transformants. The host strain for gene knockout was the *A. niger* FGSC A1279  $\Delta$ *gcnE* mutant. Gene deletion transformants were confirmed by PCR amplification using internal primers (Table S3 and Fig. S1).



### 3. Results and discussion

#### 3.1. Chemical diversity of *A. niger* FGSC A1279 $\Delta$ *gcnE* mutant

To investigate the effect of *gcnE* on *A. niger* secondary metabolism, we constructed a  $\Delta$ *gcnE* mutant and a *gcnE* complementation (*gcnE*-com) strain using homologous recombination (Fig. S1). Three types of solid media (WATM, CYA and YES) commonly used for fungal secondary metabolite production were chosen (Frisvad and Thrane, 1987; Rank et al., 2012) to investigate metabolite changes in the  $\Delta$ *gcnE* mutant and *gcnE*-com strain. Crude extract from cultures was subjected to LC/HR-ESIMS analysis. No compounds were produced in *A. niger* FGSC A1279 cultures. Metabolite profiling by LC/HR-ESIMS analysis demonstrated that multiple SMs were produced in all of three  $\Delta$ *gcnE* mutant cultures and the SM profile of *gcnE*-com strain was similar with the wild strain (Fig. 1), suggesting that *gcnE* deletion activated the synthesis of SM products in *A. niger*. It has been reported that, in *Aspergilli*, histone acetylation usually correlates with transcriptional activation and chromatin rearrangement (Baker and Grant, 2007; Reyes-Dominguez et al., 2008). The *gcnE* gene is an epigenetic regulator that is a key histone acetyltransferase in the SAGA/ADA complex. In primary metabolism, GcnE increases the acetylation level of the proline utilization cluster, as shown by chromatin immunoprecipitation in *A. nidulans* (Reyes-Dominguez et al., 2008). However, the role of GcnE might vary for different secondary metabolite gene clusters in fungi. For example, inactivation of *gcnE* abolishes the production of aflatoxin in *A. flavus* (Lan et al., 2016) and reduces the production of four known metabolites (orsellinic acid, sterigmatocystin, penicillin and terrequinone) in *A. nidulans*, suggesting that *gcnE* is required for transcription activation in these two *Aspergillus* strains (Nutzmann et al., 2011). However, recent genome-wide ChIP-seq experiments revealed that most SM cluster genes in *A. nidulans* are poorly decorated with activating heterochromatic marks (histone demethylation and acetylation), even under inducing conditions (Connolly et

al., 2013; Gacek-Matthews et al., 2016), implying that *gcnE* may not be required for transcriptional activation of certain SM gene clusters. Therefore, GcnE may have diverse effects on regulating SMs among different *Aspergilli*. This result was in accordance with the effects of histone deacetylase HdaA in *A. nidulans*. Deletion of HdaA induces the biosynthesis of sterigmatocystin, penicillin, and norsolorinic acid in *A. nidulans*, but does not change the expression of terraquinone A (Shwab et al., 2007). A similar phenomenon was found in the plant pathogen *Fusarium graminearum*. Deletion of heterochromatin protein 1 (*hep1*) in *F. graminearum* leads to the activation of the aurofusarin gene cluster while repressed the deoxynivalenol cluster (Gacek and Strauss, 2012).

### **3.2. Dereplication of newly produced compounds in *A. niger* $\Delta$ *gcnE* mutant**

To further analyze the chemical diversity in the  $\Delta$ *gcnE* mutant, we used WATM medium since more SM products are likely to be produced on this medium (Fig. 1). Dereplication using the Antibase database (H, 2012) led to the identification of 11 SM products in the *A. niger*  $\Delta$ *gcnE* mutant (Fig. 2 and Table S4), all of which were known fungal metabolites, including 9 compounds discovered in *Aspergilli*. The ion peak with  $t_R$  (7.84) min was likely to be carbonarone A, originally isolated from the fungal strain *A. carbonarius* WZ-4-11 (Zhang et al., 2007). Funalenone ( $t_R$  = 8.42 min) belongs to polyketide compound group of phenalenones which have diverse structures and biological activities (Gao et al., 2016). The biosynthetic gene cluster of funalenones has been fully characterized (Gao et al., 2016). Pestalamide A may have been eluted at  $t_R$  (10.10 min), which was isolated from fungal strain *Pestalotiopsis theae* W148 (Ding et al., 2008). The highly toxic metabolites fumonisin B2 ( $t_R$  = 10.94 min) and B4 ( $t_R$  = 11.92 min) were detected in the *A. niger*  $\Delta$ *gcnE* mutant. *A. niger* NRRL3 (ATCC9029), the starting strain of FGSC A1279, is generally regarded as safe (GRAS) in the industry and should not produce any toxins or toxic compounds. This is particularly true for *A. niger* FGSC A1279 which is used for gluconic acid production (Baker,

2006). These results indicate that deletion of *gcnE* had an epigenetic impact on its SM production. Five polyketide derivatives (dimeric naphthopyrone family), aurasperone E ( $t_R$  = 13.70 min), fonsecinone B ( $t_R$  = 14.50 min), isoaurasperone A ( $t_R$  = 14.77 min), aurasperone A ( $t_R$  = 15.66 min) and asperpyrone C ( $t_R$  = 16.33 min) were also found in the extracts of the *A. niger*  $\Delta gcnE$  mutant (Fig. 2). Taken together, these results indicated that the newly identified compounds were likely to be polyketide metabolites.

### 3.3. Structural elucidation by HR-ESIMS and 1D/2D NMR

To verify the identity of the newly emerged metabolites in the  $\Delta gcnE$  mutant, the strain was cultivated on WATM medium at 2-L scale and crude extract was subjected to chemical workup and semi-prep HPLC purification, yielding six pure compounds **1** (3.7 mg), **2** (1.8 mg), **3** (7.8 mg), **4** (9.2 mg), **5** (1.5 mg), and **6** (3.0 mg).

The molecular formula of nigerpyrone (**1**) was established as  $C_{12}H_{10}O_2$  by HR-ESIMS (187.07  $[M+H]^+$ ). Dereplication using the Antibase database suggested that this might be a new compound. The molecular formula suggested eight degrees of unsaturation. The  $^{13}C$  and HSQC NMR data of **1** in  $CD_3OD$  (Fig. S2) revealed one benzene/aromatic ring ( $\delta_C$  128.38-136.30 ppm), one carbonyl ( $\delta_C$  178.14 ppm), and four olefinic carbons ( $\delta_C$  117.36-117.52 ppm and 158.45-169.48 ppm), consistent with the presence of five double bonds and one ring system. This analysis accounted for seven double bonds, suggesting the presence of an additional ring in the structure of **1**. Analysis of the  $^1H$ - $^1H$  COSY spectrum revealed two contiguous spin systems, one consisting of H-10 through H-14, and the other consisting of H-2 to H-3 (Fig. 3 and Fig. S2). The connection of these two spin systems at C-6 was evident by the HMBC correlation of H-5 and H-8 to C-6 (Fig. 3), which was further confirmed by the correlation of H-8 to C-5 in the HMBC spectrum. The connection of methylene to the benzene ring was confirmed by the HMBC correlation of H-8 to C-9 (Fig. 3). The structure was further confirmed by comparing  $^{13}C$  experimental data with published data (Dai et al.,

2007) and predicted  $^{13}\text{C}$  NMR data (Elyashberg et al., 2010) (Fig. S2). The resulting  $r^2$  value of 0.9992 indicated that the proposed structure was correct (Bremser, 1978; Elyashberg et al., 2010). Inspection of the  $^1\text{H}$ ,  $^{13}\text{C}$ , and HR-ESIMS data of compounds **2** and **3** (Fig. S3 and S4) indicated that both **2** and **3** were known compounds (Zhang et al., 2007; Ding et al., 2008): carbonarone A (**2**) was isolated from *A. carbonarius* WZ-4-11 (Zhang et al., 2007) and pestalamide A (**3**) was isolated from *Pestalotiopsis theae* W148 (Ding et al., 2008). Comparison of  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of compounds **1**, **2** and **3** (Fig. S2, S3 and S4) confirmed the structure of the new compound **1**.

Inspection of the  $^1\text{H}$ ,  $^{13}\text{C}$  and HR-ESIMS data of compounds **4** (Fig. S5), **5** (Fig. S6) and **6** (Fig. S7), indicated that they were known. Compound **4** is produced by *A. niger* FO-5904 (Inokoshi et al., 1999), **5** by *A. niger* CMI-IMI 205879 (Priestap, 1984), and **6** by *A. fonsecaeus* NRRL 67, O 16-1 (Priestap, 1984).

### **3.4. Putative biosynthetic gene clusters of compounds from the *AgcnE* mutant**

Homologous BLAST search and comparison of the literature allowed the identification of the putative biosynthetic gene clusters of known compounds discovered from the  $\Delta gcnE$  mutant (Table S5). Genes for synthesizing funalenone (**4**) were identified in *Penicillium herquei* (Gao et al., 2016). By homologous search against the annotated genome of *A. niger* CBS513.88, we identified five homologous genes in *A. niger* responsible for the biosynthesis of funalenone (Fig. S8 and Table S5). The literature (Chiang et al., 2010) indicated that monodictyphenone might be the precursor of aurasperone E (**5**), aurasperone A (**6**) and other derivatives such as fonsecinone B, isoaurasperone A, and asperpyrone C. In *A. nidulans*, the key biosynthetic enzyme of monodictyphenone is proposed to be a non-reducing polyketide synthase (NR-PKS) *mdpG* (Chiang et al., 2010). We therefore performed a homologous search of the *A. niger* genome using *mdpG* as the reference query, and identified an open reading frame (ORF) An11g07310 located in contig 11, annotated as a NR-PKS with

multiple domains of KS-AT-PT-PP (Table S5). Other proposed biosynthetic genes involved in the biosynthesis of **5** and **6** were not located in the close proximity of An11g07310 and were scattered in other genomic loci. However, genes for secondary metabolites are generally tightly clustered.

### 3.5. Characterization of key biosynthetic genes for pestalamide A

Inspection of the structures of **1-3** led to speculation that their precursors contain one phenylacetate and two acetate units, suggesting that compounds **1-3** may be synthesized via a polyketide biosynthetic pathway. The incorporation of phenylacetate in fungal secondary metabolism is also observed in the biosynthesis of the  $\beta$ -lactam antibiotic penicillin G in *Penicillium chrysogenum* (Koetsier et al., 2009). *In vitro* assays show that the gene product PhlB is a phenylacetyl-CoA ligase responsible for the activation of phenylacetate into phenylacetyl-SCoA (Koetsier et al., 2009). We performed a homolog search of PhlB in *A. niger*, and identified ORF An09g01820, annotated as a ferulate:CoA ligase (AMP-forming) (Fig. 4A and Table S6), with moderate sequence identity (35%) to PhlB.

Analysis of genes in proximity to An09g01820 (*epaB*) identified a candidate NR-PKS gene cluster (*epa*) spanning a 23.9-kb genomic region (Fig. 4A and Table S6). The *epa* cluster possesses six ORFs (Table S6). *In silico* analysis indicated that *epaA* (An09g01860) encodes a typical Clade III NR-PKS (2,617 amino acids in length) with a domain organization of SAT-KS-AT-PT-ACP-Met-R domains, which shares high amino acid sequence identity (97%) with AzaA polyketide synthase (2,599 amino acids in length) in the biosynthesis of azaphilones from *A. niger* ATCC 1015 (Zabala et al., 2012). The gene product EpaC is annotated as an acyl-CoA transferase. Adjacent to *epaA*, there are three genes, orf1-3, encoding oxidoreductase, 3-hydroxybenzoate 4-hydroxylase, and salicylate hydroxylase (Table S6).

298 To determine if the *epa* cluster was responsible for synthesizing pestalamide A (**3**), we  
299 knocked out the PKS gene (An09g01860, *epaA*) by homologous recombination (Table S3  
300 and Fig. S1). Inactivation of the *epaA* gene abolished the production of **1-3** (Fig. 4B), and  
301 complementation strain (*epaA-com*) could produce compound **1-3** (Fig. S1 and Fig. 4B),  
302 suggesting that the identified biosynthetic genes were responsible for the synthesis of **1-3**. We  
303 also performed cell-free extract experiments, in which all biosynthetic enzymes and cofactors  
304 were available for the biosynthesis of **3**, to test if **1** and **2** were the biosynthetic precursors of  
305 **3** (data not shown). Feeding purified **1** and **2** into cell-free extracts of the  $\Delta$ *gcnE* mutant did  
306 not produce **3**, suggesting that **1** and **2** were by-products in the biosynthesis of **3**. Based on  
307 the genetic and cell-free extract experiments, a biosynthetic model for **1-3** is proposed in Fig.  
308 4C. We propose that the biosynthesis of **1-3** begins with the polyketide assembly by EpaA to  
309 form phenylacetyl triketide precursor from successive condensation of two malonyl-CoA,  
310 presumably with one phenylacetyl-CoA starter unit. For the nigerpyrone (**1**) biosynthesis, the  
311 reactive polyketide chain is released as an aldehyde (**8**) through the R-domain, in a manner  
312 similar to the previously characterized 3-methylorcinaldehyde synthase in *Acremonium*  
313 *strictum* (Bailey et al., 2007). The cyclization and dehydration of **8** may create nigerpyrone  
314 (**1**). This nonenzymatic rearrangement to form pyrone moiety from aldehyde was also  
315 observed in the plant metabolites, arabidopyl derivatives (Weng et al., 2012). For the  
316 biosynthesis of **2** and **3**, an extra methyl group will be added through the C-methyltransferase  
317 domain, followed by the reduction to generate **8a**, which undergoes oxidation and  
318 transamination to produce the intermediate **9**. The candidate gene products for this series of  
319 biotransformation could be orf1-3 (Table S6). The cyclization of **9** produces carbonarone A  
320 (**2**). We propose that EpaC (An09g01800), an acyl-CoA transferase, could catalyse the  
321 transfer of 2-methylsuccinyl-CoA, a common intermediate in the ethylmalonyl-CoA pathway  
322 (Erb et al., 2009), to generate the final product pestalamide A (**3**).

## 4. Conclusions

In conclusion, we confirmed that deletion of *gcnE* resulted in the production of 12 polyketide metabolites in an *A. niger* FGSC A1279  $\Delta gcnE$  mutant. Chemical workup of organic extracts of the culture broth from the  $\Delta gcnE$  mutant identified six pure compounds, including one new compound nigerpyrone (**1**) and five known fungal polyketide metabolites (**2-6**). The structures of these compounds were confirmed by HR-ESIMS and 1D/2D NMR. Finally, the biosynthetic gene cluster for **1-3** was verified via gene knockout and complementation, and a biosynthetic model of this group of pyran-based fungal metabolites was proposed. To our best knowledge, this was the first time to confirm that the inactivation of *gcnE* resulted in activating the biosynthesis of polyketide metabolites in *A. niger* FGSC A1279.

## Declarations of interest

None.

## Acknowledgements

This work was supported the Natural Science Foundation of Guangdong Province (grant number 2017A030313097), the Science and Technology Planning Project of Guangdong Province (grant numbers 2016A050503016 and 2016A010105004), the Science and Technology Planning Project of Guangzhou City (grant number 201510010191), the Fundamental Research Funds for the Central Universities (grant number 2015ZP032), and the China Scholarship Council (CSC) fund (grant number 201606155032).

## References

- Andersen MR, Salazar MP, Schaap PJ, van de Vondervoort PJ, Culley D, Thykaer J, Frisvad JC, Nielsen KF, Albang R, Albermann K, Berka RM, Braus GH, Braus-Stromeyer SA, Corrochano LM, Dai Z, van Dijck PW, Hofmann G, Lasure LL, Magnuson JK, Menke H, Meijer M, Meijer SL, Nielsen JB, Nielsen ML, van Ooyen AJ, Pel HJ, Poulsen L, Samson RA, Stam H, Tsang A, van den Brink JM, Atkins A, Aerts A, Shapiro H, Pangilinan J, Salamov A, Lou Y, Lindquist E, Lucas S, Grimwood J, Grigoriev IV, Kubicek CP, Martinez D, van Peij NN, Roubos JA, Nielsen J, Baker SE, 2011. Comparative genomics of citric-acid-producing *Aspergillus niger* ATCC 1015 versus enzyme-producing CBS 513.88. *Genome Res.* 21(6), 885-897.
- Arentshorst M, Lagendijk EL, Ram AF, 2015. A new vector for efficient gene targeting to the pyrG locus in *Aspergillus niger*. *Fungal Biol. Biotechnol.* (2), 2.
- Bailey AM, Cox RJ, Harley K, Lazarus CM, Simpson TJ, Skellam E, 2007. Characterisation of 3-methylorcinolaldehyde synthase (MOS) in *Acremonium strictum*: first observation of a reductive release mechanism during polyketide biosynthesis. *Chem. Commun.* (39), 4053-4055.
- Baker SE, 2006. *Aspergillus niger* genomics: past, present and into the future. *Med. Mycol.* 44 Suppl 1, S17-21.
- Baker SP, Grant PA, 2007. The SAGA continues: expanding the cellular role of a transcriptional co-activator complex. *Oncogene* 26(37), 5329-5340.
- Bayram O, Krappmann S, Ni M, Bok JW, Helmstaedt K, Valerius O, Braus-Stromeyer S, Kwon NJ, Keller NP, Yu JH, Braus GH, 2008. VelB/VeA/LaeA complex coordinates light signal with fungal development and secondary metabolism. *Science* 320(5882), 1504-1506.
- Bok JW, Hoffmeister D, Maggio-Hall LA, Murillo R, Glasner JD, Keller NP, 2006. Genomic mining for *Aspergillus* natural products. *Chem. Biol.* 13(1), 31-37.
- Brakhage AA, 2013. Regulation of fungal secondary metabolism. *Nat. Rev. Microbiol.* 11(1), 21-32.
- Bremser W, 1978. Hose-a novel substructure code. *Analytica Chimica Acta* 103(4), 355-365.
- Carvalho ND, Arentshorst M, Jin Kwon M, Meyer V, Ram AF, 2010. Expanding the ku70 toolbox for filamentous fungi: establishment of complementation vectors and recipient strains for advanced gene analyses. *Appl. Microbiol. Biotechnol.* 87(4), 1463-1473.
- Chiang YM, Meyer KM, Praseuth M, Baker SE, Bruno KS, Wang CC, 2011. Characterization of a polyketide synthase in *Aspergillus niger* whose product is a precursor for both dihydroxynaphthalene (DHN) melanin and naphtho-gamma-pyrone. *Fungal Genet. Biol.* 48(4), 430-437.
- Chiang YM, Szewczyk E, Davidson AD, Entwistle R, Keller NP, Wang CC, Oakley BR, 2010. Characterization of the *Aspergillus nidulans* monodictyphenone gene cluster. *Appl. Environ. Microbiol.* 76(7), 2067-2074.
- Connolly LR, Smith KM, Freitag M, 2013. The *Fusarium graminearum* histone H3 K27 methyltransferase KMT6 regulates development and expression of secondary metabolite gene clusters. *PLoS Genet.* 9(10), e1003916.
- Dai J, Liu Y, Jia H, Zhou YD, Nagle DG, 2007. Benzochromenones from the marine crinoid *Comantheria rotula* inhibit hypoxia-inducible factor-1 (HIF-1) in cell-based reporter assays and differentially suppress the growth of certain tumor cell lines. *J. Nat. Prod.* 70(9), 1462-1466.
- Ding G, Jiang L, Guo L, Chen X, Zhang H, Che Y, 2008. Pestalazines and pestalamides, bioactive metabolites from the plant pathogenic fungus *Pestalotiopsis theae*. *J. Nat. Prod.* 71(11), 1861-1865.
- Elyashberg M, Williams AJ, Blinov K, 2010. Structural revisions of natural products by Computer-Assisted Structure Elucidation (CASE) systems. *Nat. Prod. Rep.* 27(9), 1296-1328.
- Erb TJ, Fuchs G, Alber BE, 2009. (2S)-Methylsuccinyl-CoA dehydrogenase closes the ethylmalonyl-CoA pathway for acetyl-CoA assimilation. *Mol. Microbiol.* 73(6), 992-1008.
- Fisch KM, Gillaspay AF, Gipson M, Henrikson JC, Hoover AR, Jackson L, Najar FZ, Wagele H, Cichewicz RH, 2009. Chemical induction of silent biosynthetic pathway transcription in *Aspergillus niger*. *J. Ind. Microbiol. Biotechnol.* 36(9), 1199-1213.



Frisvad JCThrane U, 1987. Standardized high-performance liquid chromatography of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-VIS spectra (diode array detection). *J. Chromatogr.* 404(1), 195-214.

Gacek-Matthews A, Berger H, Sasaki T, Wittstein K, Gruber C, Lewis ZA, Strauss J, 2016. KdmB, a Jumonji histone H3 demethylase, regulates genome-wide H3K4 trimethylation and is required for normal induction of secondary metabolism in *Aspergillus nidulans*. *PLoS Genet.* 12(8), e1006222.

Gacek A Strauss J, 2012. The chromatin code of fungal secondary metabolite gene clusters. *Appl. Microbiol. Biotechnol.* 95(6), 1389-1404.

Gao SS, Duan A, Xu W, Yu P, Hang L, Houk KN, Tang Y, 2016. Phenalenone polyketide cyclization catalyzed by fungal polyketide synthase and flavin-dependent monooxygenase. *J. Am. Chem. Soc.* 138(12), 4249-4259.

Hiort J, Maksimenka K, Reichert M, Perovic-Ottstadt S, Lin WH, Wray V, Steube K, Schaumann K, Weber H, Proksch P, Ebel R, Muller WE, Bringmann G, 2004. New natural products from the sponge-derived fungus *Aspergillus niger*. *J. Nat. Prod.* 67(9), 1532-1543.

Inglis DO, Binkley J, Skrzypek MS, Arnaud MB, Cerqueira GC, Shah P, Wymore F, Wortman JR, Sherlock G, 2013. Comprehensive annotation of secondary metabolite biosynthetic genes and gene clusters of *Aspergillus nidulans*, *A. fumigatus*, *A. niger* and *A. oryzae*. *BMC Microbiol.* 13, 91.

Inokoshi J, Shiomi K, Masuma R, Tanaka H, Yamada H, Omura S, 1999. Funalenone, a novel collagenase inhibitor produced by *Aspergillus niger*. *J. Antibiot (Tokyo)*. 52(12), 1095-1100.

Koetsier MJ, Jekel PA, van den Berg MA, Bovenberg RA, Janssen DB, 2009. Characterization of a phenylacetate-CoA ligase from *Penicillium chrysogenum*. *Biochem. J.* 417(2), 467-476.

Laatsch H, 2012. AntiBase 2012 The natural compound identifier. Wiley-VCH Verlag GmbH & Co, KGaA.

Lan H, Sun R, Fan K, Yang K, Zhang F, Nie XY, Wang X, Zhuang Z, Wang S, 2016. The *Aspergillus flavus* histone acetyltransferase aflGcnE regulates morphogenesis, aflatoxin biosynthesis, and pathogenicity. *Front. Microbiol.* 7, 1324.

Nielsen KF, Mogensen JM, Johansen M, Larsen TO, Frisvad JC, 2009. Review of secondary metabolites and mycotoxins from the *Aspergillus niger* group. *Anal. Bioanal. Chem.* 395(5), 1225-1242.

Nutzmann HW, Reyes-Dominguez Y, Scherlach K, Schroeckh V, Horn F, Gacek A, Schumann J, Hertweck C, Strauss J, Brakhage AA, 2011. Bacteria-induced natural product formation in the fungus *Aspergillus nidulans* requires Saga/Ada-mediated histone acetylation. *Proc. Natl. Acad. Sci. USA* 108(34), 14282-14287.


Pel HJ, de Winde JH, Archer DB, Dyer PS, Hofmann G, Schaap PJ, Turner G, de Vries RP, Albang R, Albermann K, Andersen MR, Bendtsen JD, Benen JA, van den Berg M, Breestraat S, Caddick MX, Contreras R, Cornell M, Coutinho PM, Danchin EG, Debets AJ, Dekker P, van Dijck PW, van Dijk A, Dijkhuizen L, Driessen AJ, d'Enfert C, Geysens S, Goosen C, Groot GS, de Groot PW, Guillemette T, Henrissat B, Herweijer M, van den Hombergh JP, van den Hondel CA, van der Heijden RT, van der Kaaij RM, Klis FM, Kools HJ, Kubicek CP, van Kuyk PA, Lauber J, Lu X, van der Maarel MJ, Meulenberg R, Menke H, Mortimer MA, Nielsen J, Oliver SG, Olsthoorn M, Pal K, van Peij NN, Ram AF, Rinas U, Roubos JA, Sagt CM, Schmoll M, Sun J, Ussery D, Varga J, Vervecken W, van de Vondervoort PJ, Wedler H, Wosten HA, Zeng AP, van Ooyen AJ, Visser J, Stam H, 2007. Genome sequencing and analysis of the versatile cell factory *Aspergillus niger* CBS 513.88. *Nat. Biotechnol.* 25(2), 221-231.

Priestap HA, 1984. New naphthopyrones from *Aspergillus fonsecaeus*. *Tetrahedron* 40(19), 3617-3624.

Rank C, Klejnstrup ML, Petersen LM, Kildgaard S, Frisvad JC, Held Gotfredsen C, Ostenfeld Larsen T, 2012. Comparative chemistry of *Aspergillus oryzae* (RIB40) and *A. flavus* (NRRL 3357). *Metabolites* 2(1), 39-56.

- Reyes-Dominguez Y, Narendja F, Berger H, Gallmetzer A, Fernandez-Martin R, Garcia I, Scazzocchio C, Strauss J, 2008. Nucleosome positioning and histone H3 acetylation are independent processes in the *Aspergillus nidulans* prnD-prnB bidirectional promoter. Eukaryot. Cell 7(4), 656-663.
- Rossano F, Ortega De Luna L, Buommino E, Cusumano V, Losi E, Catania MR, 1999. Secondary metabolites of *Aspergillus* exert immunobiological effects on human monocytes. Res. Microbiol. 150(1):13-19.
- Sanchez JF, Somoza AD, Keller NP, Wang CC, 2012. Advances in *Aspergillus* secondary metabolite research in the post-genomic era. Nat. Prod. Rep. 29(3), 351-371.
- Serra R, Braga A, Venancio A, 2005. Mycotoxin-producing and other fungi isolated from grapes for wine production, with particular emphasis on ochratoxin A. Res. Microbiol. 156(4), 515-521.
- Shwab EK, Bok JW, Tribus M, Galehr J, Graessle S, Keller NP, 2007. Histone deacetylase activity regulates chemical diversity in *Aspergillus*. Eukaryot. Cell 6(9), 1656-1664.
- Sorensen LM, Lametsch R, Andersen MR, Nielsen PV, Frisvad JC, 2009. Proteome analysis of *Aspergillus niger*: lactate added in starch-containing medium can increase production of the mycotoxin fumonisin B2 by modifying acetyl-CoA metabolism. BMC Microbiol. 9, 255.
- Szewczyk E, Nayak T, Oakley CE, Edgerton H, Xiong Y, Taheri-Talesh N, Osmani SA, Oakley BR, 2006. Fusion PCR and gene targeting in *Aspergillus nidulans*. Nat. Protoc. 1(6), 3111-3120.
- Tanaka H, Wang P-L, Yamada Late O, Tamura T, 1966. Yellow pigments of *Aspergillus niger* and *Asp. awamori*. Agric. Biol. Chem. 30(2), 107-113.
- Varga J, Rigo K, Kocsube S, Farkas B, Pal K, 2003. Diversity of polyketide synthase gene sequences in *Aspergillus* species. Res. Microbiol. 154(8), 593-600.
- Wang B, Lv Y, Li X, Lin Y, Deng H, Pan L, 2018. Profiling of secondary metabolite gene clusters regulated by LaeA in *Aspergillus niger* FGSC A1279 based on genome sequencing and transcriptome analysis. Res. Microbiol. 169, 67-77.
- Weng JK, Li Y, Mo H, Chapple C, 2012. Assembly of an evolutionarily new pathway for alpha-pyrone biosynthesis in *Arabidopsis*. Science 337(6097), 960-964.
- Wu M-YYu J-H, 2015. Epigenetics of fungal secondary metabolism related genes. In: Zeilinger S, Martín J-F, García-Estrada C editors. Biosynthesis and molecular genetics of fungal secondary metabolites, Volume 2. Springer New York, New York, pp. 29-42.
- Yin C, Wang B, He P, Lin Y, Pan L, 2014. Genomic analysis of the aconidial and high-performance protein producer, industrially relevant *Aspergillus niger* SH2 strain. Gene 541(2), 107-114.
- Yuan XL, van der Kaaij RM, van den Hondel CA, Punt PJ, van der Maarel MJ, Dijkhuizen L, Ram AF, 2008. *Aspergillus niger* genome-wide analysis reveals a large number of novel alpha-glucan acting enzymes with unexpected expression profiles. Mol. Genet. Genomics 279(6), 545-561.
- Zabala AO, Xu W, Chooi YH, Tang Y, 2012. Discovery and characterization of a silent gene cluster that produces Azaphilones from *Aspergillus niger* ATCC 1015 reveal a hydroxylation-mediated pyran-ring formation. Chem. Biol. 19(8), 1049-1059.
- Zhang Y, Zhu T, Fang Y, Liu H, Gu Q, Zhu W, 2007. Carbonarones A and B, new bioactive gamma-pyrone and alpha-pyridone derivatives from the marine-derived fungus *Aspergillus carbonarius*. J. Antibiot (Tokyo). 60(2), 153-157.

## 492 Figures



493

494 **Figure 1. SM profile of *A. niger* FGSC A1279  $\Delta gcnE$  mutant cultivated on different**


495 **media.** Blue curve,  $\Delta gcnE$  mutant; red curve, wild strain; green curve, *gcnE*

496 complementation strain (*gcnE-com*). To collect secondary metabolites, *A. niger* strains were

497 cultivated for 7 days at 25 °C in the dark using different media. The crude extract was

498 dissolved in MeOH:Milli-Q H<sub>2</sub>O (9:1) and centrifuged at 10,000 rpm for 10 min, before

499 LC/MS analysis.



**Figure 2. SM products detected in *A. niger*  $\Delta gcnE$  mutant cultivated on WATM medium.**


*A. niger*  $\Delta gcnE$  mutant was cultivated for 7 days at 25 °C in the dark on WATM medium.

High resolution mass spectrometric data were obtained using an LTQ XL/LTQ Orbitrap

Discovery MS system coupled to a Thermo Instruments HPLC system. LC/MS data were

analyzed by Xcalibur software with Antibase2012 database. Compounds that were purified

by chemical workup and characterized by spectroscopic analysis were marked in bold.




**Figure 3. Secondary metabolites isolated and characterized by HR-ESIMS and 1D/2D**

**NMR in *A. niger*  $\Delta gcnE$  mutant.** *A. niger*  $\Delta gcnE$  mutant was cultivated on WATM medium.

$^1\text{H}$  NMR,  $^{13}\text{C}$  NMR, COSY, HSQC and HMBC spectra were recorded on a Bruker Daltonics

Advance III HD 400 NMR spectrometer, using deuterated solvent methanol-d<sub>4</sub>. Chemical shift values were reported in parts per million (ppm) with tetramethylsilane (TMS) as the external standard. When higher field was needed, NMR spectra were recorded on a Bruker Daltonics Advance III HD 600 NMR. NMR data were analyzed using MestReNova 9.0.1 software and ACD/I-Lab (<http://ilab.cds.rsc.org/?cdsrd=1>). Compound structures were drawn using ChemBioDraw software. Chemical formulas and exact masses are marked for each compound. Dereplication data are in online supplemental file (Fig. S2-S7).



**Figure 4. Characterization of the biosynthetic cluster for pestalamide A.** (A) Putative gene cluster of compounds **1-3**, which was identified by the homologous BLAST search of *P. chrysogenum* PhlB gene. (B) LC/MS detection of compounds **1-3** in gene knockout and complementation strains. Retention time for purified compounds **1-3** was determined by LC/MS. *A. niger* strains were cultivated on WATM medium. (C) Proposed model of the biosynthesis of nigerpyrone (**1**), carbonarone A (**2**) and pestalamide A (**3**).

526 **Supplemental materials**

527 **Table S1.** Strains and plasmids used in this study

528 **Table S2.** Primers used for *gcnE* deletion-cassette construction and transformant  
529 identification

530 **Table S3.** Primers used to knock out *epaA* (An09g01860) and verify mutants

531 **Table S4.** Putative compounds detected in *A. niger* FGSC A1279  $\Delta gcnE$  mutant based on  
532 LC/MS data and Antibase database

533 **Table S5.** Putative biosynthetic gene clusters for known compounds detected in the  $\Delta gcnE$   
534 mutant

535 **Table S6.** Putative biosynthetic gene cluster for pestalamide A.

536

537 **Figure S1.** Design of deletion cassettes, complementation cassettes, and PCR verification of  
538 *A. niger* mutants ( $\Delta gcnE$ ,  $\Delta gcnE\Delta epaA$ , *gcnE* complementation and *epaA* complementation).

539 **Figure S2.** LC/MS and NMR data for the novel compound nigerpyrone

540 **Figure S3.** LC/MS and NMR data of compound carbonarone A

541 **Figure S4.** LC/MS and NMR data of compound pestalamide A

542 **Figure S5.** LC/MS and NMR data of compound funalenone

543 **Figure S6.** LC/MS and NMR data of compound aurasperone E

544 **Figure S7.** LC/MS and NMR data of compound aurasperone A

545 **Figure S8.** Putative biosynthetic gene clusters of the known compounds in the  $\Delta gcnE$  mutant